

Entropy and Temperature From Black-Hole/Near-Horizon-CFT Duality

Leo Rodriguez and Tuna Yildirim

Department of Physics and Astronomy, The University of Iowa, Iowa City, IA 52242

E-mail: leo-rodriguez@uiowa.edu

Abstract. We construct a 2-dimensional CFT, in the form of a Liouville theory, in the near horizon limit of 4- and 3-dimensional black holes. The near horizon CFT assumes 2-dimensional black hole solutions first introduced by Christensen and Fulling [1] and expanded to a greater class of black holes via Robinson and Wilczek [2]. The 2-dimensional black holes admit a $\text{Diff}(S^1)$ subalgebra, which upon quantization in the horizon limit becomes Virasoro with calculable central charge. This charge and lowest Virasoro eigen-mode reproduce the correct Bekenstein-Hawking entropy of the 4- and 3-dimensional black holes via the known Cardy formula [3, 4]. Furthermore, the 2-dimensional CFT's energy momentum tensor is anomalous. However, In the horizon limit the energy momentum tensor becomes holomorphic equaling the Hawking flux of the 4- and 3-dimensional black holes. This encoding of both entropy and temperature provides a uniformity in the calculation of black hole thermodynamic and statistical quantities for the non local effective action approach.

1. Introduction

The universally accepted thermodynamic and statistical properties of black holes have provided a unique insight and test for theories of quantum gravity. Though a fully formulated quantum field theory of gravity is lacking, a multitude of candidates exists, with string theory and loop quantum gravity leading in popularity. Despite a variety of theories, any serious candid should reproduce the correct form of the Bekenstein-Hawking entropy [5]

$$S_{BH} = \frac{A}{4\hbar G} \quad (1)$$

and Hawking temperature [6, 7]

$$T_H = \frac{\hbar\kappa}{2\pi}, \quad (2)$$

where A † is the horizon area and κ the surface gravity of the black hole. The fact that these quantities depend on both $\hbar$ and G is evident of their quantum gravitational origin. For a comprehensive review of approaches to quantum gravity see [8, 9, 10].

Effective theories have had much success in reproducing (1) and (2) via analysis of anomalous energy momentum tensors of non local effective actions [11] and holographic 1- and 2-dimensional conformal field theories [9], respectively. Yet, these methods remain in separate camps and a lack of uniformity in the derivation of both temperature and entropy in one simultaneous effective theory is missing.

We will briefly review some of the pioneering work inspiring this note.

1.1. Effective Action and Hawking Temperature

Analysis of an anomalous energy momentum tensor to compute (2) was first carried out by Christensen and Fulling in [1]. Considering the most general solution to the conservation equation

$$\nabla_\mu T^\mu{}_\nu = 0, \quad (3)$$

they found that by restricting to the $r - t$ plane of a free scalar field in Schwarzschild geometry the energy momentum tensor exhibits a trace anomaly leading to the result:

$$\langle T^r_t \rangle = \frac{1}{768\pi G^2 M^2} = \frac{\pi}{12} T_H^2, \quad (4)$$

which is exactly the luminosity (Hawking flux, Hawking radiation) of the 4-dimensional black hole in units $\hbar = 1$.

A similar approach was studied in [12] where the authors determined the s -wave contribution of a scalar field to the 4-dimensional effective action for an arbitrary spherically symmetric gravitational field. Applying their results to a Schwarzschild black hole, the authors showed the energy momentum tensor of the non local effective

† $A = \int d^2x \sqrt{|\gamma|}$ where γ_{ij} is obtained by setting $r = r_+$ for constant time. In the case for *Kerr* $\sqrt{|\gamma|} = (r_+^2 + J^2) \sin \theta$, which deviates from the standard form.

action to contain the Hawking Flux. Other closely related approaches for scalar fields and 2-dimensional theories include [13, 14, 15, 16, 17]

Another method for computing Hawking Radiation first introduced by Robinson and Wilczek [2] considers a quantum chiral-scalar field theory of 2-dimensions in the near horizon limit of a static 4-dimensional black hole. A 2-dimensional chiral field theory is known to exhibit a gravitational anomaly of the form:

$$\nabla_\mu T^\mu{}_\nu = \frac{1}{96\pi\sqrt{-g^{(2)}}}\epsilon^{\gamma\rho}\partial_\rho\partial_\lambda\Gamma^{(2)\lambda}{}_{\nu\gamma}, \quad (5)$$

where $g^{(2)}{}_{\mu\nu}$ contains the leftover components of the 4-dimensional metric which are not redshifted away in the near horizon limit of the functional

$$S_{\text{free scalar}} = \frac{1}{2} \int d^4x \sqrt{-g} \nabla_\mu \varphi \nabla^\mu \varphi. \quad (6)$$

Robinson and Wilczek showed in the near horizon regime of a Schwarzschild black hole, that to ensure a unitary quantum field theory the black hole should radiate as a thermal bath of temperature equaling T_H . In other words, quantum gravitational effects in the near horizon regime cancel the chiral/gravitational anomaly [18]. This method has been expanded to include gauge/gravitational anomalies and covariant anomalies [19, 20, 21, 22] and has successfully reproduced the correct black hole temperature for charged-rotating black holes [23, 24], dS/AdS black holes [25, 26], rotating dS/AdS black holes [27, 28], black rings and black string [29, 30], 3-dimensional black holes [31, 32] and black holes of non spherical topologies [33]. This method provides a fundamental reason for black hole thermodynamics based on symmetry principles of a near horizon quantum field theory. It also provides a 2-dimensional analogue for higher dimensional black holes besides the Schwarzschild case. This is a rather useful fact since the Ricci tensor in 2-dimensions is always Einstein with proportionality constant $\frac{1}{2} \left(g^{(2)\mu\nu} R^{(2)}{}_{\mu\nu} \right) \S$. Thus classically, in 2-dimensions, there are no general relativistic dynamics and any gravitational effects that are present must have quantum gravitational origin with semi-classical metric $g^{(2)}{}_{\mu\nu}$.

1.2. Holographic CFT and Entropy

Applying the seminal work of Brown and Henneaux|| [34], Strominger showed that the Bekenstein-Hawking entropy for AdS_3 black holes arises naturally from microscopic states of an asymptotic CFT via Cardy's formula [35]. This work has been generalized to a class of rotating and dS/AdS black holes in various dimensions in both near horizon regime and asymptotic infinity by Carlip and others [36, 37, 38, 39, 40, 41]. Recently,

§ In 2-dimensions the curvature of any Riemannian manifold is completely characterized by its scalar variant. This is because any 2-form has only one independent component. Thus for any Riemannian-Levi-Cevita connection 2-form $\omega_{\alpha\beta}$, $d\omega_{12} = K \text{vol}^2$, where $K = \frac{1}{(2)((2)-1)} R^{(2)}$ is the Gauss curvature

|| The algebra of the asymptotic symmetry group of 3-dimensional gravity is Virasoro with calculable central charge.

Barnes, Vaman and Wu generalized the method to include 4-dimensional charged non-rotating black holes [42] via uplifting the 4-dimensional black holes to 6-dimensional pure gravitational solutions. Though slightly different in their approaches, the general idea is as follows: there exists a set of diffeomorphisms preserving vector fields either in the near horizon/asymptotic symmetries of the metric or boundary conditions at spacial infinity as in [40]. This set of diffeomorphisms includes a $\text{Diff}(S^1)$ subalgebra parametrized by some discrete set of vector fields ξ_n for all $n \in \mathbb{Z}$ such that:

$$i \{ \xi_m, \xi_n \} = (m - n) \xi_{m+n} \quad (7)$$

Upon quantization, Brown and Henneaux showed

$$[\xi_m, \xi_n] = (m - n) \xi_{m+n} + \frac{c}{12} m (m^2 - 1) \delta_{m+n,0} \quad (8)$$

where c is a calculable central extension. The Bekenstein-Hawking Entropy is then given by Cardy's Formula [3, 4] in terms of c and Δ_0 , where Δ_0 is the eigen-mode of ξ_0 , via:

$$S = 2\pi \sqrt{\frac{c \cdot \Delta_0}{6}} \quad (9)$$

Applying the above outline to the 2-dimensional dilaton black hole

$$\begin{aligned} ds^2 &= g_{\mu\nu}^D dx^\mu dx^\nu \\ &= - \left[(\lambda x)^2 - \frac{2M}{\lambda} (\lambda x)^3 \right] dt^2 + \left[(\lambda x)^2 - \frac{2M}{\lambda} (\lambda x)^3 \right]^{-1} dx^2, \end{aligned} \quad (10)$$

Cadoni computed the following central extension and zero mode [43]:

$$c = 48 \frac{M^2}{\lambda^2} \text{ and } \Delta_0 = \frac{M^2}{2\lambda^2} \quad (11)$$

Cadoni further showed by conformally mapping (10) to the s -wave sector of the Schwarzschild metric:

$$g_{\mu\nu}^{(2)} = 2\phi g_{\mu\nu}^D \quad (12)$$

with

$$\lambda^2 = \frac{1}{G} \quad (13)$$

and

$$x = \frac{G}{r} \quad (14)$$

(11) and (9) reproduced the correct Bekenstein-Hawking Entropy for the respective 4-dimensional black hole. In other words, together with Robinson and Wilczek's results [2] both entropy and temperature of the Schwarzschild black hole induces some semi-classical theory of spacetime

$$ds^2 = - \left(1 - \frac{2GM}{r} \right) dt^2 + \left(1 - \frac{2GM}{r} \right)^{-1} dr^2. \quad (15)$$

Our goal will be to expound and extend this idea to include a greater class of rotating, charged and other types of black holes.

1.3. *Kerr/CFT Correspondence*

A recent study by Guica, Hartman, Song and Strominger [44] proposed that the near horizon geometry of an extremal Kerr black hole is holographically dual to a 2-dimensional chiral CFT[¶] with non vanishing left central extension c_L . By constructing the Frolov-Thorne vacuum for generic Kerr geometry [45], which reduces to the Hartle-Hawking vacuum [46] for $J \rightarrow 0$, the authors obtain a non vanishing left Frolov-Thorne temperature T_L in the near horizon, extremal limit. This temperature together with c_L inside the Thermal Cardy Formula [47] reproduces the correct Bekenstein-Hawking entropy for the extremal Kerr black hole:

$$S_{BH} = \frac{\pi^2}{3} c_L T_L \quad (16)$$

This correspondence has been extended to various exotic black holes in string theory, higher dimensional theories and gauged supergravities to name a few [48, 47, 49].

One of the main arguments of the *Kerr/CFT* correspondence is to apply the rich ideas of holographic duality to more astrophysical objects/black-holes, such as the nearly extremal GRS 1915+105, a binary black hole system 11000 pc away in Aquila [50]. In [44] the authors show that GRS 1915+105 is holographically dual to a 2-dimensional chiral CFT with $c_L = (2 \pm 1) \times 10^{79}$ and in the extremal limit the inner most stable circular orbit corresponds to the horizon. Thus, the authors conclude, any radiation emanating from the inner most circular orbit should be well described by the 2-dimensional chiral CFT, making the *Kerr/CFT* correspondence an essential theoretical tool in an astrophysical observation.

Despite the various models observationalists employ they all incorporate four main quantities: black hole mass M and spin J , poloidal magnetic field at the horizon B_0 and (Eddington) luminosity L for both supermassive [51] and stellar [52] black holes. This provides a new testable playing field for holography, i.e. to use some induced 2-dimensional CFT in the near horizon regime of extremal and non-extremal black holes to model the four main quantities in accordance with observation. In particular, the origin or mechanism of B_0 is unclear from a theoretical standpoint, since it must be due to an accreting disk for non gauged black holes. Yet, it might find its origin in some black-hole/CFT duality.

The goal of this note is to model the near horizon regime via a 2-dimensional CFT as in the *Kerr/CFT* correspondence. This is done by combining the ideas from effective action approach and holographic duality and thus encodes both Hawking temperature and Bekenstein-Hawking entropy. Our final results hold in both extremal and non extremal cases.

¶ Distinct from the 2-dimensional chiral-scalar field theory employed by Robinson and Wilczek.

1.4. Outline of Main Idea

We will model the near horizon regime with a 2-dimensional Liouville type quantum field theory, which is well understood [53],

$$S_{Liouville} = \frac{1}{96\pi} \int d^2x \sqrt{-g^{(2)}} \{ -\Phi \square_{g^{(2)}} \Phi + 2\Phi R^{(2)} \}. \quad (17)$$

We make this choice based on the fact that in this regime all mass and angular terms of (6) fall off exponentially fast upon transformation from $r \rightarrow r_*$, where $\frac{\partial r}{\partial r_*} = f(r)$ [2] and $g^{(2)}_{tt} = -f(r)$. This leaves us with an infinite collection of 2-dimensional free scalars in spherically symmetric spacetime $g^{(2)}_{\mu\nu}$ ⁺. The effective action of 2-dimensional free scalars is given by the Polyakov action [54, 55]:

$$\Gamma_{Polyakov} = \frac{1}{96\pi} \int d^2x \sqrt{-g^{(2)}} R^{(2)} \frac{1}{\square_{g^{(2)}}} R^{(2)} \quad (18)$$

and integrating out Φ in $S_{Liouville}$ yields $\Gamma_{Polyakov}$. In the case where the original 4-dimensional metric is not spherically symmetric [23, 24], a $U(1)$ gauge sector appears in addition to (18), which adds a gauge anomaly to Robinson and Wilczek's method for computing Hawking Radiation. Yet in our analysis, in addition to the Hawking flux, the presence of the $U(1)$ field only contributes a $\sim \frac{e^2}{4\pi} \mathcal{A}_t^2 \Big|_{Horizon}$ to the holomorphic energy momentum tensor, which is the total flux of the gauge field. Thus it will suffice to only consider the field theory of (17) since we are interested in Hawking effects.

The energy momentum tensor for (17) is defined as:

$$\begin{aligned} \langle T_{\mu\nu} \rangle &= -\frac{2}{\sqrt{-g^{(2)}}} \frac{\delta S_{Liouville}}{\delta g^{(2)\mu\nu}} \\ &= -\frac{1}{48\pi} \left\{ \partial_\mu \Phi \partial_\nu \Phi - 2\nabla_\mu \partial_\nu \Phi + g^{(2)}_{\mu\nu} \left[2R^{(2)} - \frac{1}{2} \nabla_\alpha \Phi \nabla^\alpha \Phi \right] \right\} \end{aligned} \quad (19)$$

and the equation of motion for the auxiliary scalar Φ is:

$$\square_{g^{(2)}} \Phi = R^{(2)} \quad (20)$$

As an ansatz for the 2-dimensional metric $g^{(2)}_{\mu\nu}$, we choose the dimensional reduced spacetimes obtained via Robinson and Wilczek. Thus given a $g^{(2)}_{\mu\nu}$ we are free to solve (20) and (19) up to integration constants. Finally adopting Unruh Vacuum boundary conditions [56]

$$\begin{cases} T_{++} = 0 & r \rightarrow \infty, l \rightarrow \infty \\ T_{--} = 0 & r \rightarrow r_+ \end{cases} \quad (21)$$

where $x^\pm = t \pm r_*$ are light-cone coordinates, r_+ is the horizon radius defined as the largest real root of $f(r) = 0$ and l is the de Sitter radius, all relevant integration constants are determined. At the horizon and asymptotic infinity for $(\Lambda = \pm \frac{1}{l^2}) = 0$ and at the horizon only for $\Lambda \neq 0$, (19) will be dominated by one holomorphic component. This

⁺ $g^{(2)}_{\mu\nu}$ may always be assumed spherically symmetric since any Riemannian Space in 2-dimensions is conformally flat.

component equals the Hawking flux of the 4- and 3-dimensional black holes, which determines the Hawking temperature.

The entropy will be determined by counting the horizon microstates of $g^{(2)}_{\mu\nu}$ via the Cardy formula (9). Following the outline proposed in [38] we construct a near horizon $\text{Diff}(S^1)$ subalgebra satisfying (7) based on the isometries of $g^{(2)}_{\mu\nu}$. In the horizon limit ($\mathcal{I}^+$ boundary) the $\text{Diff}(S^1)$ subalgebra takes the form:

$$i2 \{ \xi_m^+, \xi_n^+ \} = (m - n) \xi_{m+n}^+, \quad (22)$$

where the factor 2 comes from neglecting the asymptotic infinity limit ($\mathcal{I}^-$ boundary). On the $\mathcal{I}^+$ boundary the energy momentum tensor is holomorphic given by the $\langle T_{++} \rangle$ component. Next, we define the charge on the $\mathcal{I}^+$ boundary*

$$Q_n = \frac{3A}{\pi G} \int dx^+ \langle T_{++} \rangle \xi_n^+, \quad (23)$$

where A is the horizon area of the higher dimensional black hole. The coefficient on the integral of Q_n is chosen such that in the case when the higher dimensional black hole is Schwarzschild $Q_n = \frac{1}{16\pi G} \int dx^+ \frac{12}{\pi(T_H^2)^2} \langle T_{++} \rangle \xi_n^+$, where we have normalized the units of the energy momentum tensor. For a 1-dimensional CFT with holomorphic energy momentum tensor $T(z)$ we have [57]:

$$\delta_{\xi(z)} T(z) = \xi T' + 2T \xi' + \frac{k}{24\pi} \xi''', \quad (24)$$

where k is the central extension associated with the CFT (17). In the case for 2-dimensional quantum scalar, $k = 1$ in agreement with (35). Thus, given the transformation (24) and compactifying the $\mathcal{I}^+$ boundary to a circle with period $(1/2 \cdot 1/T_H)$, where the $1/2$ takes the $\mathcal{I}^-$ boundary into account, we obtain the following charge algebra:

$$[Q_m, Q_n] = (m - n) Q_n + \frac{c}{12} m (m^2 - 1) \delta_{m+n,0}, \quad (25)$$

where c is the central extension associated with the horizon-microstates of $g^{(2)}_{\mu\nu}$. The Bekenstein-Hawking entropy of the 4- and 3-dimensional black holes is then given by Q_0 and c via (9).

Finally we compare our results to [43] by conformally mapping $g^D_{\mu\nu}$ to $g^{(2)}_{\mu\nu}$

$$g^{(2)}_{\mu\nu} = 2\phi g^D_{\mu\nu} \quad (26)$$

for some conformal factor $2\phi = (\lambda x)^{-2}$, where (11) is assumed invariant under conformal transformations [16, 58]. We should not that the dimensionally reduced spacetimes satisfy the equations of motion for 2-dimesnional dilaton gravity in the Schwarzschild case only. This is impart due to the fact that 2-dimensional dilaton gravity is a dimensionally reduced theory of the pure Einstein-Hilbert action with metric ansatz:

$$ds_{(4)}^2 = ds_{(2)}^2 + \frac{2}{\lambda^2} \phi d\Omega_{(2)}^2, \quad (27)$$

* Q_n is only conserved on the $\mathcal{I}^+$ boundary.

where $ds_{(2)}^2$ is as in (10), with no additional matter sources. Now, since the above metric ansatz does not incorporate axisymmetric solutions and the Reissner-Nordström solves Einstein-Hilbert-Maxwell theory, it is clear why their 2-dimensional counterparts do not solve the same field equations. Despite this short coming, we conjecture that the conformal map (26) still encodes the correct microstates for $g_{\mu\nu}^{(2)}$ based on comparisons of respective entropies obtained from (25) and (26) applied to the Reissner-Nordström case. This may be impart, since in 2-dimensions there are no classical general relativistic dynamics, as mentioned in the [Introduction](#), and $g_{\mu\nu}^{(2)}$ may be a dimensionally reduced effective metric of a ultraviolet complete 4-dimensional theory of gravity.

2. Examples from 4- and 3-Dimensions

We will now apply the method, outlined above, to various 4- and 3-dimensional black holes with zero and non zero cosmological constant and construct their Hawking flux, associated entropy and temperature.

2.1. Spherically Symmetric Solutions

In this class we will consider the Schwarzschild (*SS*) and Reissner-Nordström (*RNS*) black holes. Their 2-dimensional analogues have the form [59, 2]

$$g_{\mu\nu}^{(2)} = \begin{pmatrix} -f(r) & 0 \\ 0 & \frac{1}{f(r)} \end{pmatrix}, \quad (28)$$

where

$$f_{SS}(r) = 1 - \frac{2GM}{r} \quad (29)$$

and

$$f_{RNS}(r) = 1 - \frac{2GM}{r} + \frac{Q^2 G}{r^2} \quad (30)$$

Next, using the above ansatz and solving (20) we get:

$$\Phi_{SS} = C_2 t + C_1 r + \ln r - (1 - 2GMC_1) \ln(r - 2GM) + C_3 \quad (31)$$

and

$$\begin{aligned} \Phi_{RNS} = & C_2 t + C_1 r + \frac{C_1 \sqrt{G} (2GM^2 - Q^2)}{\sqrt{GM^2 - Q^2}} \arctan \left(\frac{GM - r}{\sqrt{G^2 M^2 - GQ^2}} \right) \\ & + 2 \ln r - (1 - GMC_1) \ln(r^2 - 2GMr + GQ^2) + C_3 \end{aligned} \quad (32)$$

Using these auxiliary fields in (19) and transforming to light cone coordinates we obtain:

$$\begin{aligned} \langle T_{++}^{SS} \rangle &= \frac{-r^4(C_1 + C_2)^2 + 8rGM - 12G^2M^2}{192\pi r^4} \\ \langle T_{--}^{SS} \rangle &= \frac{-r^4(C_1 - C_2)^2 + 8rGM - 12G^2M^2}{192\pi r^4} \\ \langle T_{+-}^{SS} \rangle &= \langle T_{-+}^{SS} \rangle = \frac{GM(r - 2GM)}{24\pi r^4} \end{aligned} \quad (33)$$

and

$$\begin{aligned}
\langle T_{++}^{RNS} \rangle &= [-r^6(C_1 + C_2)^2 + 8r^3GM - 12r^2G(GM^2 + Q^2) \\
&\quad + 24rG^2MQ^2 - 8G^2Q^4] / [192\pi a^6] \\
\langle T_{--}^{RNS} \rangle &= [-r^6(C_1 - C_2)^2 + 8r^3GM - 12r^2G(GM^2 + Q^2) \\
&\quad + 24rG^2MQ^2 - 8G^2Q^4] / [192\pi a^6] \\
\langle T_{+-}^{RNS} \rangle &= \langle T_{-+}^{RNS} \rangle = \frac{G(2rM - 3Q^2)(r^2 - 2rGM + GQ^2)}{48\pi r^6}
\end{aligned} \tag{34}$$

The fact that both energy momentum tensors are not holomorphic/anti-holomorphic signals the existence of a conformal anomaly taking the form

$$\langle T_\mu{}^\mu \rangle = -\frac{1}{24\pi} R^{(2)} \tag{35}$$

due to the trace anomaly [57]. Imposing (21) to eliminate C_1 and C_2 our final steps are to analyze $\langle T_{\mu\nu} \rangle$ at the horizon and construct the conformal map (26). At the horizon the energy momentum tensors are dominated by one holomorphic component $\langle T_{++} \rangle$ given by

$$\langle T_{++}^{SS} \rangle = \frac{1}{768\pi G^2 M^2} = \frac{\pi}{12} (T_H)^2 \tag{36}$$

and

$$\begin{aligned}
\langle T_{++}^{RNS} \rangle &= \frac{G^2(GM^2 - Q^2) \left(2M\sqrt{G(GM^2 - Q^2)} + 2GM^2 - Q^2 \right)}{48\pi \left(\sqrt{G(GM^2 - Q^2)} + GM \right)^6} \\
&= \frac{\pi}{12} (T_H)^2,
\end{aligned} \tag{37}$$

which are in agreement with Hawking's original results [6, 7].

Following [38], we compute the near horizon diffeomorphisms satisfying (7). We get:

$$\xi_n^{SS} = 4GM e^{i\kappa t} \partial_t + \frac{in(r - 2GM)^2 e^{i\kappa t}}{GM} \partial_r \tag{38}$$

and

$$\begin{aligned}
\xi_n^{RNS} &= \frac{\left(\sqrt{G(GM^2 - Q^2)} + GM \right)^3 e^{i\kappa t}}{G \left(M\sqrt{G(GM^2 - Q^2)} + GM^2 - Q^2 \right)} \partial_t + \\
&\quad \frac{in e^{i\kappa t} (r^2 - 2rGM + GQ^2)^2}{rG(rM - Q^2)} \partial_r,
\end{aligned} \tag{39}$$

where κ is the horizon surface gravity. Transforming to $x^\pm$ coordinates and taking the horizon limit, we obtain the $\mathcal{I}^+$ boundary charge algebras:

$$[Q_m, Q_n]_{SS} = (m - n)Q_n + 2GM^2 m(m^2 - 1) \delta_{m+n,0}, \tag{40}$$

and

$$\begin{aligned}
[Q_m, Q_n]_{RNS} &= (m - n)Q_n + \\
&\quad \left(M\sqrt{G(GM^2 - Q^2)} + GM^2 - \frac{Q^2}{2} \right) m(m^2 - 1) \delta_{m+n,0},
\end{aligned} \tag{41}$$

which imply

$$Q_0^{SS} = GM^2 \quad (42)$$

$$c^{SS} = 24GM^2 \quad (43)$$

and

$$Q_0^{RNS} = \frac{\left(\sqrt{G(GM^2 - Q^2)} + GM\right)^2}{4G} \quad (44)$$

$$c^{RNS} = \frac{6\left(\sqrt{G^2M^2 - GQ^2} + GM\right)^2}{G} \quad (45)$$

and using these values in (9) we get:

$$S_{SS} = 4\pi GM^2 = \frac{A}{4G} \quad (46)$$

and

$$S_{RNS} = \frac{\pi\left(\sqrt{G(GM^2 - Q^2)} + GM\right)^2}{G} = \frac{A}{4G}. \quad (47)$$

Finally, conformally mapping $g^{(2)}_{\mu\nu}$ to $g^{(D)}_{\mu\nu}$ implies:

$$\lambda_{SS}^2 = \frac{1}{G} \quad (48)$$

$$x_{SS} = \frac{G}{r} \quad (49)$$

$$c_{SS} = 48GM^2 \quad (50)$$

$$\Delta_0^{SS} = \frac{GM^2}{2} \quad (51)$$

as in [43] and

$$\lambda_{RNS}^2 = \frac{4GM^2}{\left(\sqrt{G^2M^2 - GQ^2} + GM\right)^2} \quad (52)$$

$$x_{RNS} = \frac{G(2rM - Q^2)}{2r^2M} \quad (53)$$

$$c_{RNS} = \frac{12\left(\sqrt{G^2M^2 - GQ^2} + GM\right)^2}{G} \quad (54)$$

$$\Delta_0^{RNS} = \frac{\left(\sqrt{G^2M^2 - GQ^2} + GM\right)^2}{8G} \quad (55)$$

where λ_{RNS} and x_{RNS} are such that

$$\lim_{Q \rightarrow 0} \lambda_{RNS} = \lambda_{SS} \text{ and } \lim_{Q \rightarrow 0} x_{RNS} = x_{SS} \quad (56)$$

Using (9) we find the respective entropies:

$$S_{SS} = 4\pi GM^2 = \frac{A}{4G} \quad (57)$$

and

$$S_{RNS} = \frac{\pi \left(\sqrt{G(GM^2 - Q^2)} + GM \right)^2}{G} = \frac{A}{4G}. \quad (58)$$

We see that our central extension and zero-mode relate to Cadoni's via

$$c = \frac{c_c}{2} \quad (59)$$

and

$$Q_0 = 2\Delta_0. \quad (60)$$

Yet, their respective products are equal and produce entropies in agreement with the Bekenstein-Hawking area law [5] for $\hbar = 1$. Thus in accord with our conjecture at the end of Section 1.4, we choose to conformally map into Cadoni's solution for all $g^{(2)}_{\mu\nu}$ for calculational simplicity.

We will proceed to solidify our main argument by applying the methods of Section 1.4 to several more black hole solutions of various types.

2.2. Axisymmetric Solutions

For this class we analyze the Kerr (K) and Kerr-Newman (KN) Black holes with 2-dimensional analogues [23, 59]

$$g^{(2)}_{\mu\nu} = \begin{pmatrix} -f(r) & 0 \\ 0 & \frac{1}{f(r)} \end{pmatrix}, \quad (61)$$

where

$$f(r) = \frac{\Delta}{r^2 + J^2} \quad (62)$$

and

$$\Delta = \begin{cases} r^2 - 2rGM + J^2 & K \\ r^2 - 2rGM + GQ^2 + J^2 & KN \end{cases}. \quad (63)$$

The auxiliary scalars read:

$$\begin{aligned} \Phi_K = & rC_1 + tC_2 + (C_1GM - 1) \log(r^2 - 2rGM + J^2) + \log(r^2 + J^2) + \\ & \frac{2C_1G^2M^2 \arctan\left(\frac{r-GM}{\sqrt{J^2 - G^2M^2}}\right)}{\sqrt{J^2 - G^2M^2}} + C_3 \end{aligned} \quad (64)$$

and

$$\begin{aligned} \Phi_{KN} = & rA + tC_2 + (C_1GM - 1) \log(r^2 - 2rGM + GQ^2 + J^2) + \\ & \log(r^2 + J^2) + \frac{C_1G(2GM^2 - Q^2) \arctan\left(\frac{r-GM}{\sqrt{G(Q^2 - GM^2) + J^2}}\right)}{\sqrt{G(Q^2 - GM^2) + J^2}} + C_3 \end{aligned} \quad (65)$$

from which we obtain the energy momentum tensors:

$$\begin{aligned}
\langle T_{++}^K \rangle &= - \left[r^8 (C_1 + C_2)^2 + 4r^6 J^2 (C_1 + C_2)^2 - 8r^5 GM + 6r^4 (C_1^2 J^4 + \right. \\
&\quad \left. 2C_1 J^4 C_2 + 2G^2 M^2 + J^4 C_2^2) + 16r^3 G J^2 M + 4r^2 (C_1^2 J^6 + \right. \\
&\quad \left. 2C_1 J^6 C_2 - 10G^2 J^2 M^2 + J^6 C_2^2) + 24r G J^4 M + C_2^2 J^8 + \right. \\
&\quad \left. 2C_1 J^8 C_2 - 4G^2 J^4 M^2 + J^8 C_2^2 \right] / \left[192\pi (r^2 + J^2)^4 \right] \\
\langle T_{--}^K \rangle &= - \left[r^8 (C_1 - C_2)^2 + 4r^6 J^2 (C_1 - C_2)^2 - 8r^5 GM + 6r^4 (C_1^2 J^4 - \right. \\
&\quad \left. 2C_1 J^4 C_2 + 2G^2 M^2 + J^4 C_2^2) + 16r^3 G J^2 M + 4r^2 (C_1^2 J^6 - \right. \\
&\quad \left. 2C_1 J^6 C_2 - 10G^2 J^2 M^2 + J^6 C_2^2) + 24r G J^4 M + C_1^2 J^8 - \right. \\
&\quad \left. 2C_1 J^8 C_2 - 4G^2 J^4 M^2 + J^8 C_2^2 \right] / \left[192\pi (r^2 + J^2)^4 \right] \quad (66) \\
\langle T_{+-}^K \rangle &= \langle T_{-+}^K \rangle = \frac{rGM (r^2 - 3J^2) (r^2 - 2rGM + J^2)}{24\pi (r^2 + J^2)^4}
\end{aligned}$$

and

$$\begin{aligned}
\langle T_{++}^{KN} \rangle &= - \left[r^8 (C_1 + C_2)^2 + 4r^6 J^2 (C_1 + C_2)^2 - 8r^5 GM + 6r^4 (C_1^2 J^4 + \right. \\
&\quad \left. 2C_1 J^4 C_2 + 2G^2 M^2 + 2GQ^2 + J^4 C_2^2) - 8r^3 GM (3GQ^2 - \right. \\
&\quad \left. 2J^2) + 4r^2 (C_1^2 J^6 + 2C_1 J^6 C_2 + 2G^2 (Q^4 - 5J^2 M^2) + \right. \\
&\quad \left. 2GJ^2 Q^2 + J^6 C_2^2) + 24r G J^2 M (GQ^2 + J^2) + C_1^2 J^8 + \right. \\
&\quad \left. 2C_1 J^8 C_2 - 4G^2 J^4 M^2 - 4G^2 J^2 Q^4 - 4GJ^4 Q^2 + J^8 C_2^2 \right] / \\
&\quad \left[192\pi (r^2 + J^2)^4 \right] \\
\langle T_{--}^{KN} \rangle &= - \left[r^8 (C_1 - C_2)^2 + 4r^6 J^2 (C_1 - C_2)^2 - 8r^5 GM + 6r^4 (C_1^2 J^4 - \right. \\
&\quad \left. 2C_1 J^4 C_2 + 2G^2 M^2 + 2GQ^2 + J^4 C_2^2) - 8r^3 GM (3GQ^2 - \right. \\
&\quad \left. 2J^2) + 4r^2 (C_1^2 J^6 - 2C_1 J^6 C_2 + 2G^2 (Q^4 - 5J^2 M^2) + \right. \\
&\quad \left. 2GJ^2 Q^2 + J^6 C_2^2) + 24r G J^2 M (GQ^2 + J^2) + C_1^2 J^8 - \right. \\
&\quad \left. 2C_1 J^8 C_2 - 4G^2 J^4 M^2 - 4G^2 J^2 Q^4 - 4GJ^4 Q^2 + J^8 C_2^2 \right] / \\
&\quad \left[192\pi (r^2 + J^2)^4 \right] \quad (67) \\
\langle T_{+-}^{KN} \rangle &= \langle T_{-+}^{KN} \rangle = \left[G (2r^3 M - 3r^2 Q^2 - 6r J^2 M + J^2 Q^2) (r^2 - \right. \\
&\quad \left. 2rGM + GQ^2 + J^2) \right] / \left[48\pi (r^2 + J^2)^4 \right]
\end{aligned}$$

which exhibits conformal anomaly (35). Applying (21) and taking the horizon limit we obtain the holomorphic pieces

$$\langle T_{++}^K \rangle = \frac{\pi (G^2 M^2 - J^2)}{12 (4\pi GM \sqrt{G^2 M^2 - J^2} + 4\pi G^2 M^2)^2} = \frac{\pi}{12} (T_H)^2 \quad (68)$$

and

$$\begin{aligned}
\langle T_{++}^{KN} \rangle &= - \frac{G (Q^2 - GM^2) + J^2}{48\pi \left(\left(\sqrt{G^2 M^2 - GQ^2 - J^2} + GM \right)^2 + J^2 \right)^2} \\
&= \frac{\pi}{12} (T_H)^2 \quad (69)
\end{aligned}$$

agreeing with Hawking's result [6, 7]. Next, from (26) and applying similar boundary conditions as in (56) we obtain

$$\lambda_K^2 = \frac{4GM^2}{(\sqrt{G^2M^2 - J^2} + GM)^2 + J^2} \quad (70)$$

$$x_K = \frac{rG}{r^2 + J^2} \quad (71)$$

$$c_K = \frac{12 \left((\sqrt{G^2M^2 - J^2} + GM)^2 + J^2 \right)}{G} \quad (72)$$

$$\Delta_0^K = \frac{(\sqrt{G^2M^2 - J^2} + GM)^2 + J^2}{8G} \quad (73)$$

and

$$\lambda_{KN}^2 = \frac{4GM^2}{\left(\sqrt{G^2M^2 - GQ^2 - J^2} + GM \right)^2 + J^2} \quad (74)$$

$$x_{KN} = \frac{2aGM - GQ^2}{2a^2M + 2J^2M} \quad (75)$$

$$c_{KN} = \frac{12 \left(\left(\sqrt{G^2M^2 - GQ^2 - J^2} + GM \right)^2 + J^2 \right)}{G} \quad (76)$$

$$\Delta_0^{KN} = \frac{\left(\sqrt{G^2M^2 - GQ^2 - J^2} + GM \right)^2 + J^2}{8G} \quad (77)$$

which give the respective entropies:

$$S_K = 2\pi M \left(\sqrt{G^2M^2 - J^2} + GM \right) = \frac{A}{4G} \quad (78)$$

and

$$S_{KN} = \pi \left(2M \left(\sqrt{G(GM^2 - Q^2) - J^2} + GM \right) - Q^2 \right) = \frac{A}{4G} \quad (79)$$

reproducing the Bekenstein-Hawking area law [5] and continuing the trend of Section 2.1.

2.3. Spherically Symmetric SSdS and Rotating BTZ

Now, we turn our attention to black holes with non zero cosmological constant:

$$\Lambda = \begin{cases} \frac{1}{l^2} & dS \\ -\frac{1}{l^2} & AdS \end{cases}, \quad (80)$$

where l is the de Sitter radius. In this black hole class we consider the spherically symmetric dS (SSdS) with line element

$$ds^2 = - \left(1 - \frac{2GM}{r} - \frac{r^2\Lambda}{3} \right) dt^2 + \left(1 - \frac{2GM}{r} - \frac{r^2\Lambda}{3} \right)^{-1} dr^2 + r^2 d\Omega \quad (81)$$

and the 3-dimensional *BTZ* black hole with line element

$$ds^2 = - \left(-8GM + \frac{r^2}{l^2} + \frac{16GJ^2}{r^2} \right) dt^2 + \left(-8GM + \frac{r^2}{l^2} + \frac{16GJ^2}{r^2} \right)^{-1} dr^2 + r^2 \left(d\phi - \frac{4GJ}{r^2} dt \right)^2. \quad (82)$$

Their 2-dimensional analogues [32, 25] are as in (61) where

$$f(r) = \begin{cases} 1 - \frac{2GM}{r} - \frac{r^2\Lambda}{3} & SSdS \\ -8GM + \frac{r^2}{l^2} + \frac{16GJ^2}{r^2} & BTZ \end{cases} \quad (83)$$

Following the steps outlined in Section 1.4 we obtain the energy momentum tensors:

$$\begin{aligned} \langle T_{++}^{SSdS} \rangle &= - \left[r^4 (3C_1^2 + 6C_1C_2 + 3C_2^2 - 4\Lambda) + 24r^3GM\Lambda - 24rGM + 36G^2M^2 \right] / [576\pi r^4] \\ \langle T_{--}^{SSdS} \rangle &= \left[r^4 (-3C_1^2 + 6C_1C_2 - 3C_2^2 + 4\Lambda) - 24r^3GM\Lambda + 24rGM - 36G^2M^2 \right] / [576\pi r^4] \\ \langle T_{+-}^{SSdS} \rangle &= \langle T_{-+}^{SSdS} \rangle = - \left[r^6\Lambda^2 - 3r^4\Lambda + 12r^3GM\Lambda - 18rGM + 36G^2M^2 \right] / [432\pi a^4] \end{aligned} \quad (84)$$

and

$$\begin{aligned} \langle T_{++}^{BTZ} \rangle &= - \left[r^6 (C_1^2 l^2 + 2C_1 l^2 C_2 - 32GM + l^2 C_2^2) + 384r^4 G^2 J^2 - 1536r^2 G^3 J^2 l^2 M + 2048G^4 J^4 l^2 \right] / [192\pi r^6 l^2] \\ \langle T_{--}^{BTZ} \rangle &= - \left[r^6 (C_1^2 l^2 - 2C_1 l^2 C_2 - 32GM + l^2 C_2^2) + 384r^4 G^2 J^2 - 1536r^2 G^3 J^2 l^2 M + 2048G^4 J^4 l^2 \right] / [192\pi r^6 l^2] \\ \langle T_{+-}^{BTZ} \rangle &= \langle T_{-+}^{BTZ} \rangle = - \left[(r^4 + 48G^2 J^2 l^2) (r^4 - 8r^2 G l^2 M + 16G^2 J^2 l^2) \right] / [48\pi r^6 l^4] \end{aligned} \quad (85)$$

with conformal anomaly (35). Applying (21) we obtain the holomorphic piece

$$\langle T_{++} \rangle = \frac{\pi}{12} (T_H)^2 \quad (86)$$

for both spacetimes in their respective horizon limits and agreeing as before with Hawking's results [6, 7]. Next, their respective entropies are computed via (9), (26),

$$\begin{aligned} \lambda_{SSdS}^2 &= - \left[16GM^2\Lambda^2 \left(\sqrt{\Lambda^3 (9G^2M^2\Lambda - 1)} - 3GM\Lambda^2 \right)^{2/3} \right] / \\ &\quad \left[\left((\sqrt{3} - i) \left(\sqrt{\Lambda^3 (9G^2M^2\Lambda - 1)} - 3GM\Lambda^2 \right)^{2/3} + (-\sqrt{3} - i) \Lambda \right)^2 \right] \end{aligned} \quad (87)$$

$$x_{SSdS} = \frac{r^3\Lambda + 6GM}{6rM} \quad (88)$$

$$c_{SSdS} = - \left[3 \left((\sqrt{3} - i) \left(\sqrt{\Lambda^3 (9G^2M^2\Lambda - 1)} - 3GM\Lambda^2 \right)^{2/3} \right) \right]$$

$$+ \left(-\sqrt{3} - i \right) \Lambda \Big)^2 \Big] / \left[G \Lambda^2 \left(\sqrt{\Lambda^3 (9G^2 M^2 \Lambda - 1)} - 3GM \Lambda^2 \right)^{2/3} \right] \quad (89)$$

$$\Delta_0^{SSdS} = - \left[\left(\left(\sqrt{3} - i \right) \left(\sqrt{\Lambda^3 (9G^2 M^2 \Lambda - 1)} - 3GM \Lambda^2 \right)^{2/3} + \left(-\sqrt{3} - i \right) \Lambda \right)^2 \right] / \left[32G \Lambda^2 \left(\sqrt{\Lambda^3 (9G^2 M^2 \Lambda - 1)} - 3GM \Lambda^2 \right)^{2/3} \right] \quad (90)$$

and

$$\lambda_{BTZ}^2 = \frac{4GM^2}{\sqrt{\sqrt{G^2 l^2 (l^2 M^2 - J^2)} + Gl^2 M}} \quad (91)$$

$$x_{BTZ} = \frac{(-r^4 + r^2 l^2 - 16G^2 J^2 l^2 + 8r^2 Gl^2 M)}{(2r^2 l^2 M)} \quad (92)$$

$$c_{BTZ} = \frac{12 \sqrt{\sqrt{G^2 l^2 (l^2 M^2 - J^2)} + Gl^2 M}}{G} \quad (93)$$

$$\Delta_0^{BTZ} = \frac{\sqrt{\sqrt{G^2 l^2 (l^2 M^2 - J^2)} + Gl^2 M}}{8G} \quad (94)$$

reproducing the Bekenstein-Hawking area law [5]

$$S = \frac{A}{4G} \quad (95)$$

in both cases via (9). Thus by modeling the near horizon regime with a 2-dimensional CFT, we have computed both entropy and temperature of 4-dimensional Schwarzschild, Reissner-Nordström, Kerr, Kerr-Newman, Spherically Symmetric dS and of 3-dimensional BTZ black hole.

3. Conclusion

To conclude, we have analyzed quantum black hole properties via non local effective action in the near horizon regime. For a relatively large class of black holes, including dS and AdS solutions in 3- and 4-dimensions, both entropy and temperature are computed from 2-dimensional conformal field theory techniques in the near horizon regime. The 2-dimensional CFT was modeled via a Liouville action with 2-dimensional black hole solutions given by Robinson and Wilczek's dimensional reduction first discussed in [2]. These 2-dimensional black holes exhibit a $\text{Diff}(S^1)$ subalgebra, up to conformal transformation, first discovered by Cadoni [43] for the s -wave sector of a Schwarzschild black hole. Analysis of the anomalous energy momentum tensor of the Liouville theory and the $\text{Diff}(S^1)$ subalgebra reproduces the Hawking temperature and Bekenstein-Hawking entropy for the respective 4- and 3-dimensional black holes.

The anomalous contribution (35) signals interesting physics as Christensen and Fulling showed [1]. In this case the anomaly relates to quantum black hole physics and in fact for Schwarzschild

$$\frac{1}{16} \langle T_{\mu}^{\mu} \rangle = -\frac{\pi}{12} (T_H)^2, \quad (96)$$

which is not the case for any other spacetime considered above except in their respective holomorphic limits. Though the 2-dimensional trace anomaly did not factor much into the main calculations of this note, it remains an interesting feature to interpret especially for the non Schwarzschild cases.

The methods outlined in Section 1.4 seem to be universal at least in 3- and 4-dimensions. It remains interesting, for future work, to generalize (23) and (25) to general black holes/strings in arbitrary dimensions as Peng, Wu [30] and Xu, Chen [28] have done for Robinson and Wilczek's gauge gravitational anomaly cancellation method.

Acknowledgments

We thank Vincent Rodgers for suggesting this problem and support. We thank Steven Carlip for enlightening discussions, comments and encouragement. We thank Edwin Barnes, Philip Kaaret, Diana Vaman and Leopoldo Pando Zayas for enlightening discussions. We also thank the entire [Diffeomorphisms and Geometry Research Group](#) of the University of Iowa for continued support, helpful discussions and comments.

We would also like to thank the editorial staff of *Class. Quantum Grav.* and the referees for helping us correct our work and pointing us to interesting and useful literature.

-
- [1] S M Christensen and S A Fulling. Trace anomalies and the hawking effect. *Physics Review D*, 15(8):2088–2104, 1977.
 - [2] S P Robinson and F Wilczek. Relationship between hawking radiation and gravitational anomalies. *Phys. Rev. Lett.*, 95, 2005.
 - [3] H W J Blöte, J A Cardy, and M P Nightingale. *Phys. Rev. Lett.*, 56(742), 1986.
 - [4] J A Cardy. *Nucl. Phys.*, B270(186), 1986.
 - [5] Jacob D Bekenstein. Black holes and entropy. *Phys. Rev. D*, 7:2333 – 2346, 1973.
 - [6] S W Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43:199 – 220, 1975.
 - [7] S W Hawking, J M Bardeen, and B Carter. The four laws of black hole mechanics. *Communications in Mathematical Physics*, 31:161 – 170, 1973.
 - [8] G K AU. The quest for quantum gravity. [arXiv:gr-qc/9506001v1](#).
 - [9] S Carlip. Horizon constraints and black hole entropy. [arXiv:gr-qc/0508071v1](#).
 - [10] Carlo Rovelli. Strings, loops and others: a critical survey of the present approaches to quantum gravity. [arXiv:gr-qc/9803024v3](#).
 - [11] A Wipf. Quantum fields near black holes. [arXiv:hep-th/9801025v1](#).
 - [12] V Mukhanov and A Wipf. On 4d-hawking radiation from effective action. *Physics Letters B*, 332(3-4):282–291, 1994. [arXiv:hep-th/9403018v1](#).

- [13] R. Balbinot and A. Fabbri. 4d quantum black hole physics from 2d models? *Physics Letters B*, 459(1-3):112–118, 1999. [arXiv:gr-qc/9904034v2](#).
- [14] R. Balbinot and A. Fabbri. Hawking radiation by effective two-dimensional theories. *Phys. Rev. D*, 59(4), 1999. [arXiv:hep-th/9807123v1](#).
- [15] M Cadoni. Trace anomaly and hawking effect in generic 2d dilaton gravity theories. *Nuclear Physics B*, 57(1-3):188–191, 1996. [arXiv:gr-qc/9510012v1](#).
- [16] M Cadoni. Trace anomaly and hawking effect in 2d dilaton gravity theories. [arXiv:gr-qc/9612041v1](#).
- [17] Shuang-Qing Wu, Jun-Jin Peng, and Zhan-Yue Zhao. Anomalies, effective action and hawking temperatures of a anomalies, effective action and hawking temperatures of a schwarzschild black hole in the isotropic coordinates. *Class. Quantum Grav.*, 25(135001), 2008. [arXiv:0803.1338v5](#).
- [18] S Das, S P. Robinson, and E C. Vagenas. Gravitational anomalies: A recipe for hawking radiation. *International Journal of Modern Physics D*, 17(3-4):533–539, 2008.
- [19] R Banerjee. Covariant anomalies, horizons and hawking radiation. *Int. J. Mod. Phys.*, D17:2539–2542, 2009. [arXiv:0807.4637v1](#).
- [20] R Banerjee and S Kulkarni. Hawking radiation and covariant anomalies. *Phys. Rev. D*, 77(2), 2008. [arXiv:0707.2449v3](#).
- [21] R Banerjee and S Kulkarni. Hawking radiation, effective actions and covariant boundary conditions. *Phys. Lett. B*, 659(4):827–831, 2008. [arXiv:0709.3916v3](#).
- [22] R Banerjee and S Kulkarni. Hawking radiation, covariant boundary conditions and vacuum states. *Phys. Rev. D*, 79(8), 2009. [arXiv:0810.5683v2](#).
- [23] S Iso, H Umetsu, and F Wilczek. Anomalies, hawking radiations and regularity in rotating black holes. *Phys. Rev. D*, 74(4), 2006. [arXiv:hep-th/0606018v3](#).
- [24] K Murata and J Soda. Hawking radiations from rotating black holes and gravitational anomalies. *Phys. Rev. D*, 74(4), 2006. [arXiv:hep-th/0606069v2](#).
- [25] S Gangopadhyay. Hawking radiation from black holes in de sitter spaces via covariant anomalies. *Gen Relativ Gravit*, 2009. DOI 10.1007/s10714-009-0900-0.
- [26] Q-Q Jiang. Hawking radiation from black holes in de sitter spaces. *Class. Quantum Grav.*, 24:4391–4406, 2007.
- [27] Q-Q Jiang and S-Q Wu. Hawking radiation from rotating black holes in anti-de sitter spaces via gauge and gravitational anomalies. *Phys. Lett. B*, 647:200–206, 2007.
- [28] Z Xu and B Chen. Hawking radiation from general kerr-(anti)de sitter black holes. *Phys. Rev. D*, 75(2), 2007. [arXiv:hep-th/0612261v3](#).
- [29] B Chen and W He. Hawking radiation of black rings from anomalies. *Class. Quantum Grav.*, 25(13), 2008. [arXiv:0705.2984v2](#).
- [30] J-J Peng and S-Qing Wu. Covariant anomalies and hawking radiation from charged rotating black strings in anti-de sitter spacetimes. *Physics Letters B*, 661:300–306, 2008.
- [31] S Nam and J-D Park. Hawking radiation from covariant anomalies in (2+1)-dimensional black holes. *Class. Quantum Grav.*, 26:15pp, 2009.
- [32] M.R. Setare. Gauge and gravitational anomalies and hawking radiation of rotating btz black holes. *The Euro. Phys. Journal C - Particles and Fields*, 49(3):865–868, 2006. [arXiv:hep-th/0608080v1](#).
- [33] E Papantonopoulos and P Skamagoulis. Hawking radiation via gravitational anomalies in non-spherical topologies. *Phys. Rev. D*, 79(8), 2009. [arXiv:0812.1759v3](#).
- [34] J D Brown and M Henneaux. *J. Math. Phys.*, 27:489, 1986.
- [35] A Strominger. Black hole entropy from near-horizon microstates. *JHEP*, 2(9), 1998. [arXiv:hep-th/9712251v3](#).
- [36] S Carlip. Black hole entropy from conformal field theory in any dimension. *Phys. Rev. Lett.*, 82(14):2828–2831, 1998. [arXiv:hep-th/9812013v3](#).
- [37] S Carlip. Conformal field theory, (2+1)-dimensional gravity, and the btz black hole. *Class. Quantum Grav.*, 22(12):R85–R123, 2005. [arXiv:gr-qc/0503022v4](#).

- [38] S Carlip. Entropy from conformal field theory at killing horizons. *Class. Quantum Grav.*, 16(10):3327–3348, 1999. [arXiv:gr-qc/9906126v2](#).
- [39] G Kanga, J-I Kogab, and M-I Park. Near-horizon conformal symmetry and black hole entropy in any dimension. *Phys. Rev. D*, 70(2), 2004. [arXiv:hep-th/0402113v1](#).
- [40] S Silva. Black hole entropy and thermodynamics from symmetries. *Class. Quantum Grav.*, 19(15):3947–3961, 2002. [arXiv:hep-th/0204179](#).
- [41] Rabin Banerjee, Sunandan Gangopadhyay, and Shailesh Kulkarni. Hawking radiation and near horizon universality of chiral virasoro algebra. [arXiv:0804.3492v2](#).
- [42] E Barnes, D Vaman, and C Wu. All 4-dimensional static, spherically symmetric, 2-charge abelian kaluza-klein black holes and their cft duals. [arXiv:0908.2425v2](#).
- [43] M Cadoni. Statistical entropy of the schwarzschild black hole. *Modern Physics Letters A*, 21(24):1879–1887, 2006.
- [44] M Guica, T Hartman, W Song, and A Strominger. The kerr/cft correspondence. *Phys. Rev. D*, 80(12), 2009. [arXiv:0809.4266](#).
- [45] V P Frolov and K S Thorne. Renormalized stress-energy tensor near the horizon of a slowly evolving, rotating black hole. *Phys. Rev. D*, 39(8):2125–2154, 1989.
- [46] J B Hartle and S W Hawking. Path-integral derivation of black-hole radiance. *Phys. Rev. D*, 13(8):2188–2203, 1976.
- [47] D D. K. Chow, M. Cvetič, H. Lü, and C. N. Pope. Extremal black hole/cft correspondence in (gauged) supergravities. *Phys. Rev. D*, 79(8), 2009. [arXiv:0812.2918](#).
- [48] T Azeanagi, N Ogawa, and S Terashima. The kerr/cft correspondence and string theory. *Phys. Rev. D*, 79(10), 2009.
- [49] H. Lü, J Mei, and C. N. Pope. Kerr-ads/cft correspondence in diverse dimensions. *JHEP*, (4):54, 2009. [arXiv:0811.2225](#).
- [50] J E McClintock, R Shafee, R Narayan, and R A Remillard. The spin of the near-extreme kerr black hole grs 1915+105. *The Astrophysical Journal*, 652(1):518–539, 2006. [arXiv:astro-ph/0606076](#).
- [51] R A Daly. Bounds on black hole spins. *The Astrophysical Journal*, 696(1):L32–L36, 2009.
- [52] R A Remillard and J E McClintock. X-ray properties of black-hole binaries. *Annual Review of Astronomy and Astrophysics*, 44:49–92, 2006. [arXiv:astro-ph/0606352](#).
- [53] N Seiberg. Notes on quantum liouville theory and quantum gravity. *Progr. of Theo. Phys. Supp.*, (102):319–349, 1990.
- [54] V Mukhanov and S Winitzki. *Intro. To Quantum Effects In Gravity*. Cambridge, 2007.
- [55] A M Polyakov. Quantum geometry of bosonic string. *Phys. Lett. B*, 103:207, 1981.
- [56] W G Unruh. Notes on black-hole evaporation. *Physics Review D*, 14(4):870–892, 1976.
- [57] P D Francesco, P Mathieu, and D Sénéchal. *Conformal Field Theory*. Springer, 1997.
- [58] M Cadoni and S Mignemi. On the conformal equivalence between 2d black holes and rindler spacetime. *Physics Letters B*, 358(3-4):217–222, 1995. [arXiv:gr-qc/9505032v1](#).
- [59] S Iso, H Umetsu, and F Wilczek. Hawking radiation from charged black holes via gauge and gravitational anomalies. *Phys. Rev. Lett.*, 96(15), 2006. [arXiv:hep-th/0602146v3](#).